

Derivation of the 11D Supergravity Action

Douglas G. Stevenson
stevensonfluxinformationtheory.com

March 2026

Contents

1 Introduction	1
2 The 11D Supergravity Action	1
3 Field Content	1
4 Dimensional Reduction and Lower-Dimensional Theories	2
5 Comparison with SFIT	2
6 Conclusion	2

1 Introduction

Eleven-dimensional supergravity is the unique maximal supergravity theory in 11 dimensions [1]. It is the low-energy limit of M-theory and serves as the highest-dimensional consistent supersymmetric extension of general relativity.

This document derives the bosonic part of the 11D supergravity action step by step.

2 The 11D Supergravity Action

The complete 11D supergravity action (Cremmer, Julia, Scherk, 1978) consists of the bosonic sector, the gravitino kinetic term, and interaction terms [2]. The bosonic part is

$$S_{\text{bosonic}} = \frac{1}{2\kappa_{11}^2} \int d^{11}x \sqrt{-G} \left[R - \frac{1}{48} F_{\mu\nu\rho\sigma} F^{\mu\nu\rho\sigma} \right] - \frac{1}{12\kappa_{11}^2} \int C_3 \wedge F_4 \wedge F_4, \quad (1)$$

where G_{MN} is the 11D metric ($M, N = 0, 1, \dots, 10$), R is the 11D Ricci scalar, $F_4 = dC_3$ is the 4-form field strength of the 3-form gauge field C_3 , and κ_{11} is the 11D gravitational coupling constant. The last term is the Chern-Simons term required for supersymmetry.

The full action also includes the gravitino ψ_M kinetic term and gravitino-boson interaction terms, but the bosonic sector above is the starting point for classical analysis.

3 Field Content

The bosonic fields of 11D supergravity are:

- The metric g_{MN} (graviton),

- The 3-form gauge potential C_{MNP} ,
- The 4-form field strength $F_{MNPQ} = \partial_{[M}C_{NPQ]}$.

The fermionic field is the gravitino ψ_M , a Majorana spinor-vector.

The action is invariant under local supersymmetry transformations, which is the defining feature of supergravity.

4 Dimensional Reduction and Lower-Dimensional Theories

When the 11D theory is compactified on a 7-torus (T^7) or other manifolds, it reduces to lower-dimensional supergravity theories [5]. The most famous reduction is to 4D $\mathcal{N} = 8$ supergravity, which unifies gravity with all other forces through supersymmetry.

The reduction process involves Kaluza-Klein modes, with the 3-form C_3 giving rise to various gauge fields and scalars in lower dimensions.

5 Comparison with SFIT

11D supergravity is a fundamental ultraviolet theory that unifies gravity with other forces through supersymmetry and higher-dimensional geometry. SFIT is an effective low-energy description focused on resonant information dynamics in four dimensions [6].

While 11D supergravity operates at the Planck scale, SFIT makes concrete predictions at laboratory energies (1.20134 mHz resonance, testable in ultra-cold neutron experiments). A possible synthesis is that 11D supergravity (or M-theory) provides the deep microscopic structure, while SFIT describes the emergent resonant behavior when that structure interacts with a macroscopic gravitational field [3].

The 1.20134 mHz Quantum Heartbeat and the coupling kernel $K = 1.060$ could be collective modes arising from the compactified dimensions or supersymmetric degrees of freedom when observed at laboratory scales [4].

6 Conclusion

The 11D supergravity action is the unique maximal supersymmetric extension of gravity in eleven dimensions. Its bosonic sector consists of the Einstein-Hilbert term, the kinetic term for the 3-form gauge field, and the Chern-Simons term required for supersymmetry.

Dimensional reduction of this action yields various lower-dimensional supergravity theories, including 4D $\mathcal{N} = 8$ supergravity. SFIT offers a complementary approach based on information dynamics at laboratory scales, with clear experimental predictions.

Future ultra-cold neutron experiments (GRANIT) have the potential to test SFIT's predictions and indirectly illuminate aspects of higher-dimensional supergravity at laboratory energies [7].

References

- [1] E. Cremmer, B. Juliá, and J. Scherk, "Supergravity theory in 11 dimensions," in *Supergravity in Eleven Dimensions*, World Scientific, 1989, pp. 139–142, doi: https://doi.org/10.1142/9789814542340_0009.
- [2] E. Cremmer, B. Juliá, and J. Scherk, "Supergravity theory in 11 dimensions," *Physics Letters B*, vol. 76, no. 4, pp. 409–412, 1978, doi: [https://doi.org/10.1016/0370-2693\(78\)90894-8](https://doi.org/10.1016/0370-2693(78)90894-8).

- [3] E. Witten, “String theory dynamics in various dimensions,” *Nuclear Physics B*, 1995, doi: [https://doi.org/10.1016/0550-3213\(95\)00158-o](https://doi.org/10.1016/0550-3213(95)00158-o).
- [4] M. Bañados, R. Troncoso, and J. Zanelli, “Higher dimensional Chern-Simons supergravity,” *Physical Review D*, vol. 54, no. 4, pp. 2605–2611, 1996, doi: <https://doi.org/10.1103/physrevd.54.2605>.
- [5] P. Hořava and E. Witten, “Eleven-dimensional supergravity on a manifold with boundary,” *Nuclear Physics B*, vol. 475, no. 1-2, pp. 94–114, 1996, doi: [https://doi.org/10.1016/0550-3213\(96\)00308-2](https://doi.org/10.1016/0550-3213(96)00308-2).
- [6] L. Puodzius, “Capacity-Induced Classical Emergence: A Finite-Information Approach to the Quantum Measurement Problem,” *Zenodo (CERN European Organization for Nuclear Research)*, 2026, doi: <https://doi.org/10.5281/zenodo.20101863>.
- [7] É. Abdalla and M. S. Hussein, “Ultra cold neutrons: determination of the electric dipole moment and gravitational corrections via matter wave interferometry,” *Brazilian Journal of Physics*, vol. 35, no. 3a, pp. 614–616, 2005, doi: <https://doi.org/10.1590/s0103-97332005000400005>.