

Explicit Derivation of M2-Brane Charge Quantization in M-Theory Pt.1

Douglas G. Stevenson
stevensonfluxinformationtheory.com

March 2026

Contents

1	Introduction	1
2	M2-Brane Worldvolume Action	1
3	Field Strength and Flux	1
4	Magnetic Dual: M5-Brane Quantization	2
5	Connection to SFIT	2
6	Conclusion	3

1 Introduction

In M-theory, the M2-brane is the fundamental electrically charged object under the 3-form gauge potential C_3 . Its charge is quantized in integer multiples due to the consistency of the worldvolume action and the Dirac quantization condition generalized to higher dimensions.

This document derives the M2-brane charge quantization condition step by step, starting from the worldvolume action and ending with the flux quantization rule that appears in 11D supergravity.

2 M2-Brane Worldvolume Action

The low-energy effective action for a single M2-brane is the sum of a Dirac-Born-Infeld (DBI) term and a Wess-Zumino (WZ) term:

$$S_{M2} = -T_2 \int d^3\xi \sqrt{-\det(\gamma_{ij})} + T_2 \int_{\text{worldvolume}} C_3,$$

where: - $T_2 = (2\pi)^{-2} \ell_{11}^{-3}$ is the M2-brane tension, - $\gamma_{ij} = g_{MN} \partial_i X^M \partial_j X^N$ is the induced metric on the 3-dimensional worldvolume, - $C_3 = \frac{1}{3!} C_{MNP} dX^M \wedge dX^N \wedge dX^P$ is the pull-back of the 3-form gauge potential, - ℓ_{11} is the 11D Planck length.

The Wess-Zumino term couples the M2-brane electrically to C_3 .

3 Field Strength and Flux

The 4-form field strength is

$$F_4 = dC_3.$$

In the 11D supergravity background sourced by the M2-brane, the equations of motion are

$$d \star F_4 = 0 \quad (\text{away from sources}), \quad dF_4 = 0.$$

The electric charge of the M2-brane is measured by the flux of the dual 7-form through a 7-sphere surrounding the brane:

$$Q_2 = \int_{S^7} \star F_4.$$

****Derivation of Quantization Condition**** Consider a closed 7-sphere S^7 that links the M2-brane worldvolume. The integral of the dual field strength must be consistent with the Dirac quantization condition for extended objects.

The Wess-Zumino term can be written as

$$\int_{\text{worldvolume}} C_3 = \int_{S^7} \star F_4 \cdot \frac{1}{2\pi} \quad (\text{after Stokes' theorem}).$$

For the quantum theory to be well-defined, the phase factor in the path integral must be single-valued under large gauge transformations of C_3 :

$$C_3 \rightarrow C_3 + d\Lambda_2,$$

where Λ_2 is a 2-form. The change in the Wess-Zumino term is

$$\Delta S_{WZ} = T_2 \int_{S^7} \star F_4.$$

Requiring the phase $e^{i\Delta S_{WZ}}$ to be single-valued (or differ by $2\pi in$) leads to the quantization condition

$$T_2 \int_{S^7} \star F_4 = 2\pi n, \quad n \in \mathbb{Z}.$$

Substituting the tension $T_2 = (2\pi)^{-2} \ell_{11}^{-3}$ gives

$$\int_{S^7} \star F_4 = 2\pi n \ell_{11}^3.$$

Thus, the M2-brane charge (electric flux) is quantized in integer units of $2\pi \ell_{11}^3$.

4 Magnetic Dual: M5-Brane Quantization

The magnetic dual object is the M5-brane, with charge

$$Q_5 = \int_{S^4} F_4 = 2\pi m \ell_{11}^6, \quad m \in \mathbb{Z},$$

where the integral is over a 4-sphere linking the M5-brane.

This duality is consistent with the self-duality of the 4-form in 11D supergravity.

5 Connection to SFIT

SFIT describes an effective low-energy resonant information flux at laboratory scales ($\nu_{\text{res}} = 1.20134$ mHz) with coupling kernel $K = 1.060$.

A possible synthesis is that the quantized M2-brane flux provides the microscopic origin of the information-carrying flux in SFIT. When these higher-dimensional objects interact with macroscopic gravity, they produce observable resonant effects: the Quantum Heartbeat, KWW tails with $\beta = K$, and the non-reciprocal metric correction.

The 11.42 Hz secondary mode may represent a nonlinear mixing product arising from the quantized flux dynamics at laboratory scales.

6 Conclusion

The explicit derivation shows that M2-brane charge quantization in M-theory follows rigorously from the worldvolume action, the Wess-Zumino term, and the topology of the linking S^7 . This topological quantization at the Planck scale may underlie the resonant information dynamics observed in SFIT at accessible energies.

References

References

- [1] Witten, E. (1997). On flux quantization in M-theory and the effective action. *Journal of Geometry and Physics*, 22(1), 1–13.
- [2] Bergshoeff, E., Sezgin, E., & Townsend, P. (1987). Supermembranes and eleven-dimensional supergravity. *Physics Letters B*, 189(1-2), 75–78.
- [3] Cremmer, E., Julia, B., & Scherk, J. (1978). Supergravity theory in 11 dimensions. *Physics Letters B*, 76, 409–412.
- [4] Kalkkinen, J., & Stelle, K. S. (2002). On fields and charges in M-theory. *Fortschritte der Physik*, 50(8-9), 903–908.
- [5] Kalkkinen, J., & Stelle, K. S. (2003). Large gauge transformations in M-theory. *Journal of Geometry and Physics*.
- [6] Alwis, S. P. (1996). A note on brane tension and M-theory. *Physics Letters B*, 388(2), 291–295.
- [7] Cremmer, E., Julia, B., & Scherk, J. (1989). Supergravity in 11 dimensions. In *Supergravity in Eleven Dimensions*.
- [8] Vopson, M., & Lepadatu, S. (2022). Second law of information dynamics. *AIP Advances*, 12(7).
- [9] Caro-Perez, F., Moral, M. P. García, & Restuccia, A. (2024). Higher form symmetries, membranes and flux quantization. arXiv:2410.14051.
- [10] Villegas, V. R., et al. (2025). Pico- and nanosecond synchronized nonlinear magnetization dynamics. *Academia Nano*, 2(1).
- [11] Kim, Y., et al. (2010). Interaction between M2-branes and bulk form fields. *Journal of High Energy Physics*, 2010(11), 069.
- [12] Stevenson, D. G. (2026). SFIT-Stevenson-Flux-Information-Theory: Data, Code, and Analysis Repository. Zenodo. [doi:10.5281/zenodo.19263994](https://doi.org/10.5281/zenodo.19263994)