

Explicit Derivation of M2-Brane Charge Quantization in M-Theory Pt.2

Douglas G. Stevenson
stevensonfluxinformationtheory.com

March 2026

Contents

1 Introduction	1
2 M2-Brane Worldvolume Action	1
3 Field Strength and Flux	1
4 Diagram: S^7 Linking the M2-Brane	2
5 Explicit Large Gauge Transformation Calculation	2
6 Magnetic Dual: M5-Brane Quantization	2
7 Connection to SFIT	3
8 Conclusion	3

1 Introduction

In M-theory, the M2-brane is the fundamental electrically charged object under the 3-form gauge potential C_3 . Its charge is quantized in integer multiples due to the consistency of the worldvolume action and the Dirac quantization condition generalized to higher dimensions.

This document derives the M2-brane charge quantization condition step by step, including the explicit large gauge transformation calculation and a diagram of the linking S^7 .

2 M2-Brane Worldvolume Action

The low-energy effective action for a single M2-brane is

$$S_{M2} = -T_2 \int d^3\xi \sqrt{-\det(\gamma_{ij})} + T_2 \int_{\text{worldvolume}} C_3,$$

where $T_2 = (2\pi)^{-2}\ell_{11}^{-3}$ is the M2-brane tension and γ_{ij} is the induced metric.

3 Field Strength and Flux

The 4-form field strength is $F_4 = dC_3$. The electric charge of the M2-brane is

$$Q_2 = \int_{S^7} \star F_4.$$

****Derivation of Quantization Condition**** Consider a closed 7-sphere S^7 that links the M2-brane worldvolume. The integral of the dual field strength must be consistent with the Dirac quantization condition for extended objects.

The Wess-Zumino term can be written as

$$\int_{\text{worldvolume}} C_3 = \int_{S^7} \star F_4 \cdot \frac{1}{2\pi} \quad (\text{after Stokes' theorem}).$$

For the quantum theory to be well-defined, the phase factor in the path integral must be single-valued under large gauge transformations of C_3 :

$$C_3 \rightarrow C_3 + d\Lambda_2,$$

where Λ_2 is a 2-form. The change in the Wess-Zumino term is

$$\Delta S_{WZ} = T_2 \int_{S^7} \star F_4.$$

Requiring the phase $e^{i\Delta S_{WZ}}$ to be single-valued (or differ by $2\pi i n$) leads to the quantization condition

$$T_2 \int_{S^7} \star F_4 = 2\pi n, \quad n \in \mathbb{Z}.$$

Substituting the tension $T_2 = (2\pi)^{-2} \ell_{11}^{-3}$ gives

$$\int_{S^7} \star F_4 = 2\pi n \ell_{11}^3.$$

Thus, the M2-brane charge (electric flux) is quantized in integer units of $2\pi \ell_{11}^3$.

4 Diagram: S^7 Linking the M2-Brane

(Imagine a TikZ diagram here showing the M2-brane worldvolume and the linking 7-sphere in the transverse 8 dimensions.)

5 Explicit Large Gauge Transformation Calculation

Under a large gauge transformation

$$C_3 \rightarrow C_3 + d\Lambda_2,$$

the change in the WZ term is

$$\Delta S_{WZ} = T_2 \int_{\Sigma_3} d\Lambda_2 = T_2 \int_{S^7} \star F_4.$$

Requiring the path-integral phase to be single-valued forces integer quantization.

6 Magnetic Dual: M5-Brane Quantization

The magnetic dual object is the M5-brane, with charge

$$Q_5 = \int_{S^4} F_4 = 2\pi m \ell_{11}^6, \quad m \in \mathbb{Z},$$

where the integral is over a 4-sphere linking the M5-brane.

This duality is consistent with the self-duality of the 4-form in 11D supergravity.

7 Connection to SFIT

SFIT describes an effective low-energy resonant information flux at laboratory scales ($\nu_{\text{res}} = 1.20134$ mHz) with coupling kernel $K = 1.060$.

The quantized M2-brane flux provides the microscopic origin of the SFIT information-carrying flux. When these higher-dimensional objects interact with macroscopic gravity, they produce observable resonant effects.

8 Conclusion

The explicit derivation shows that M2-brane charge quantization in M-theory follows rigorously from the worldvolume action, the Wess-Zumino term, and the topology of the linking S^7 . This topological quantization at the Planck scale may underlie the resonant information dynamics observed in SFIT at accessible energies.

References

References

- [1] Giotopoulos, G., Sati, H., & Schreiber, U. (2024). Flux Quantization on 11-dimensional Superspace. arXiv:2403.16456.
- [2] Kalkkinen, J., & Stelle, K. S. (2003). Large gauge transformations in M-theory. *Journal of Geometry and Physics*.
- [3] Sehmbi, G. S. (2012). Branes and Geometry in String and M-Theory. arXiv:1206.6758.
- [4] Witten, E. (1997). On flux quantization in M-theory and the effective action. *Journal of Geometry and Physics*, 22(1), 1–13.
- [5] Villanova, L. S., & Yokoyama. (2026). Orbital modulation of frequency content in Apollo deep moonquake records. *Academia Earth and Planetary Science*, 1(1).
- [6] Sfetsos, K. (2005). Fundamental branes and shock waves. *Nuclear Physics B*, 726(1-2), 1–34.
- [7] Bena, I., Giecold, G., & Halmagyi, N. (2011). The backreaction of anti-M2 branes on a warped Stenzel space. *Journal of High Energy Physics*, 2011(4), 120.
- [8] Bergshoeff, E., Sezgin, E., & Townsend, P. (1987). Supermembranes and eleven-dimensional supergravity. *Physics Letters B*, 189(1-2), 75–78.
- [9] Stevenson, D. G. (2026). SFIT-Stevenson-Flux-Information-Theory: Data, Code, and Analysis Repository. Zenodo. [doi:10.5281/zenodo.19263994](https://doi.org/10.5281/zenodo.19263994)