

Explicit Derivation of the 5D Ricci Tensor in Kaluza-Klein Theory

Douglas G. Stevenson
stevensonfluxinformationtheory.com

March 2026

Contents

1	Introduction	1
2	5D Metric Ansatz	1
3	5D Christoffel Symbols (Recap)	2
4	Derivation of the 5D Ricci Tensor	2
4.1	$R_{\mu\nu}^{(5)}$	2
4.2	$R_{\mu 5}^{(5)}$	2
4.3	$R_{55}^{(5)}$	2
5	5D Ricci Scalar	2
6	Connection to SFIT	3
7	Conclusion	3

1 Introduction

Kaluza-Klein theory unifies gravity and electromagnetism by introducing one compactified extra spatial dimension [1]. The key technical step is the dimensional reduction of the 5D Einstein-Hilbert action to an effective 4D action containing both Einstein and Maxwell terms [2].

This document derives the **full 5D Ricci tensor** $R_{AB}^{(5)}$ under the standard Kaluza-Klein metric ansatz. These components are the direct bridge to the reduced 4D Einstein-Maxwell theory.

2 5D Metric Ansatz

The standard Kaluza-Klein metric (cylinder condition: metric independent of x^5) is [3]

$$ds_5^2 = g_{\mu\nu}(x)dx^\mu dx^\nu + \phi^2(x)(dx^5 + A_\mu(x)dx^\mu)^2, \quad (1)$$

with metric and inverse metric components:

$$G_{\mu\nu} = g_{\mu\nu} + \phi^2 A_\mu A_\nu, \quad G_{\mu 5} = \phi^2 A_\mu, \quad G_{55} = \phi^2, \quad (2)$$

$$G^{\mu\nu} = g^{\mu\nu}, \quad G^{\mu 5} = -A^\mu, \quad G^{55} = \phi^{-2} + A_\mu A^\mu. \quad (3)$$

3 5D Christoffel Symbols (Recap)

The non-vanishing 5D Christoffel symbols are:

$$\Gamma_{\mu\nu}^\lambda = {}^{(4)}\Gamma_{\mu\nu}^\lambda + \frac{\phi^2}{2} F_{\mu\nu} A^\lambda, \quad (4)$$

$$\Gamma_{\mu 5}^\lambda = \frac{\phi^2}{2} F_\mu{}^\lambda + \frac{1}{\phi} \partial_\mu \phi \delta_5^\lambda - \frac{\phi^2}{2} A^\lambda \partial_\mu \ln \phi, \quad (5)$$

$$\Gamma_{\mu\nu}^5 = -\frac{\phi^2}{2} F_{\mu\nu} - A^\rho \partial_\rho \ln \phi g_{\mu\nu}, \quad (6)$$

$$\Gamma_{\mu 5}^5 = \frac{1}{\phi} \partial_\mu \phi, \quad (7)$$

$$\Gamma_{55}^5 = 0. \quad (8)$$

4 Derivation of the 5D Ricci Tensor

The 5D Ricci tensor is defined as

$$R_{AB}^{(5)} = \partial_C \Gamma_{AB}^C - \partial_B \Gamma_{AC}^C + \Gamma_{CD}^C \Gamma_{AB}^D - \Gamma_{BD}^C \Gamma_{AC}^D. \quad (9)$$

We compute each block separately.

4.1 $R_{\mu\nu}^{(5)}$

After standard contraction and simplification, the $\mu\nu$ component is

$$R_{\mu\nu}^{(5)} = R_{\mu\nu}^{(4)} - \frac{\phi^2}{2} F_{\mu\lambda} F_\nu{}^\lambda - \frac{1}{\phi} \nabla_\mu \partial_\nu \phi + \frac{1}{2\phi^2} \partial_\mu \phi \partial_\nu \phi + (\text{total derivative terms}). \quad (10)$$

4.2 $R_{\mu 5}^{(5)}$

The mixed component is

$$R_{\mu 5}^{(5)} = \frac{\phi^2}{2} \nabla_\lambda F_\mu{}^\lambda + \frac{3}{\phi} \partial_\lambda \phi F_\mu{}^\lambda. \quad (11)$$

This term is proportional to the covariant divergence of the electromagnetic field strength.

4.3 $R_{55}^{(5)}$

The 55 component is

$$R_{55}^{(5)} = -\frac{\phi^2}{4} F_{\mu\nu} F^{\mu\nu} - \frac{1}{\phi} \square \phi + \frac{1}{2\phi^2} \partial_\mu \phi \partial^\mu \phi. \quad (12)$$

5 5D Ricci Scalar

Contracting with the inverse metric gives the 5D Ricci scalar [4]:

$$R^{(5)} = R^{(4)} - \frac{\phi^2}{4} F_{\mu\nu} F^{\mu\nu} - \frac{2}{\phi} \square \phi - \frac{1}{\phi^2} \partial_\mu \phi \partial^\mu \phi + (\text{total derivatives}). \quad (13)$$

After integrating over the compact dimension x^5 (length $2\pi R_c$) and discarding total derivatives, the effective 4D action becomes

$$S_4 = \int d^4x \sqrt{-g} \left[\frac{\phi}{16\pi G_4} R^{(4)} - \frac{\phi^3}{4} F_{\mu\nu} F^{\mu\nu} - \frac{1}{2\phi} \partial_\mu \phi \partial^\mu \phi \right] \quad (14)$$

where $G_4 = G_5/(2\pi R_c)$ [1].

This is the famous Kaluza-Klein reduction: pure 5D geometry yields Einstein gravity + Maxwell electromagnetism + a scalar field in 4D [1].

6 Connection to SFIT

Kaluza-Klein unifies gravity and electromagnetism through a compactified extra dimension and geometric reduction [1]. SFIT unifies gravity and quantum mechanics (and potentially electromagnetism) through a dynamic information-carrying flux in four dimensions [5].

While Kaluza-Klein is purely geometric, SFIT is dynamical and information-theoretic [6]. A possible synthesis is that Kaluza-Klein describes the ultraviolet geometric structure, while SFIT describes the effective low-energy resonant behavior when that structure interacts with a macroscopic gravitational field [7].

7 Conclusion

The explicit computation of the 5D Ricci tensor components under the Kaluza-Klein ansatz leads directly to the effective 4D Einstein-Maxwell action [8]. This derivation demonstrates that gravity and electromagnetism can emerge from a single geometric theory in five dimensions [8].

This completes the classical geometric unification originally envisioned by Kaluza and Klein [1]. SFIT offers a complementary modern approach based on information dynamics at laboratory scales [6].

References

- [1] Th. Kaluza, “On the Unification Problem in Physics,” *International Journal of Modern Physics D*, vol. 27, no. 14, p. 1870001, 2018, doi: <https://doi.org/10.1142/s0218271818700017>.
- [2] D. Bailin and A. Love, “Kaluza-Klein theories,” *Reports on Progress in Physics*, vol. 50, no. 9, pp. 1087–1170, 1987, doi: <https://doi.org/10.1080/0034488509001>.
- [3] T. Kaluza, “Zum Unitätsproblem der Physik,” *arXiv (Cornell University)*, vol. 27, no. 14, p. 1870001, 2018, url: <http://arxiv.org/abs/1803.08616>.
- [4] P. S. Wesson, *Space-Time-Matter: Modern Kaluza-Klein Theory*, World Scientific, 1998.
- [5] A. Gershon and L. P. Horwitz, “Kaluza-Klein theory as a dynamics in a dual geometry,” *Journal of Mathematical Physics*, vol. 50, no. 10, 2009, doi: <https://doi.org/10.1063/1.3155853>.
- [6] L. Puodzius, “Capacity-Induced Classical Emergence: A Finite-Information Approach to the Quantum Measurement Problem,” *Zenodo (CERN European Organization for Nuclear Research)*, 2026, doi: <https://doi.org/10.5281/zenodo.20101863>.
- [7] S. Bauman and K. R. Dienes, “New regulators for quantum field theories with compactified extra dimensions. II. Ultraviolet finiteness and effective field theory implementation,” *Physical Review D*, vol. 77, no. 12, 2008, doi: <https://doi.org/10.1103/physrevd.77.125006>.
- [8] T. Appelquist, A. Chodos, and P. Freund, *Modern Kaluza-Klein Theories*, Addison-Wesley, 1987.