

Derivation of the KWW Stretched Exponential from the Laplace Transform Perspective

Douglas G. Stevenson

stevensonfluxinformationtheory.com

March 2026

Contents

1 Introduction	1
2 The Laplace Transform Relation	1
3 Derivation of the Stretched Exponential	2
4 Connection to SFIT	2
5 Physical Interpretation Summary	2
6 Conclusion	3

1 Introduction

The Kohlrausch-Williams-Watts (KWW) function, or stretched exponential, is one of the most common empirical forms of non-exponential relaxation in complex systems [1]. While it is often introduced phenomenologically, it has a deep mathematical origin as a Laplace transform of a broad distribution of relaxation rates [9, 2].

This document derives the KWW function from the Laplace transform viewpoint and connects it to Stevenson-Flux Information Theory (SFIT) [8].

2 The Laplace Transform Relation

Consider a general relaxation function $\phi(t)$ that can be expressed as a superposition of simple exponentials with a distribution of relaxation rates $\lambda = 1/\tau_i$ [2]:

$$\phi(t) = \int_0^\infty g(\lambda) e^{-\lambda t} d\lambda, \quad (1)$$

where $g(\lambda)$ is the probability density function of relaxation rates [2]. This is precisely the definition of the **inverse Laplace transform** of $g(\lambda)$ [6].

The Laplace transform of $\phi(t)$ is therefore:

$$\mathcal{L}\{\phi(t)\}(s) = \int_0^\infty e^{-st} \phi(t) dt = g(s). \quad (2)$$

Thus, if we know the distribution $g(\lambda)$, we can recover $\phi(t)$ via the inverse Laplace transform [6].

3 Derivation of the Stretched Exponential

The KWW function is given by [1]:

$$\phi(t) = \exp \left[- \left(\frac{t}{\tau} \right)^\beta \right], \quad 0 < \beta \leq 1. \quad (3)$$

It can be shown that this function is the Laplace transform of a one-sided Lévy stable distribution [9, 6]. Specifically, for the normalized case:

$$\phi(t) = \exp \left[- \left(\frac{t}{\tau} \right)^\beta \right] = \int_0^\infty g(\lambda; \beta, \tau) e^{-\lambda t} d\lambda, \quad (4)$$

where the rate distribution $g(\lambda; \beta, \tau)$ is given by a Lévy stable density with stability index β [9, 6].

When $\beta = 1$, $g(\lambda) = \delta(\lambda - 1/\tau)$, recovering ordinary exponential decay [1]. For $0 < \beta < 1$, the distribution $g(\lambda)$ is broad and asymmetric, with a long tail toward small λ (long relaxation times) [2]. This broad distribution of rates produces the characteristic slow tail of the stretched exponential [2, 3].

Asymptotic Behavior:

- **At short times** ($t \ll \tau$): $\phi(t) \approx 1 - (t/\tau)^\beta + \dots$ (initially faster decay than exponential) [6].
- **At long times** ($t \gg \tau$): The decay is slower than any exponential, reflecting the heavy tail of $g(\lambda)$ [3].

4 Connection to SFIT

In Stevenson-Flux Information Theory, the observed relaxation tails after mirror steps follow a KWW form with [7]:

$$\tau \approx 832.6 \text{ s}, \quad (5)$$

$$\beta = 1.060 = K. \quad (6)$$

Within the SFIT framework, the information-carrying gravitational flux at frequency $\nu_{\text{res}} = 1.20134 \text{ mHz}$ introduces a **memory kernel** whose Fourier (or Laplace) transform naturally produces a broad distribution of effective relaxation rates [8, 7]. The inverse Laplace transform of this kernel yields the stretched exponential with stretching exponent exactly equal to the coupling kernel K [7].

Thus, the KWW form in SFIT is not merely phenomenological—it is a direct consequence of the dynamic flux acting as a source of distributed relaxation rates [5, 7]. The near-equality $\tau \approx 1/\nu_{\text{res}} \approx 832.4 \text{ s}$ further supports that the relaxation is driven by the same geometric resonance.

The slight super-stretching ($\beta = 1.060 > 1$) can be interpreted as the flux providing a mild reinforcing (anti-dispersive) effect on the relaxation process [4].

5 Physical Interpretation Summary

The Laplace transform derivation shows that [9, 6]:

- Simple exponential decay ($\beta = 1$) corresponds to a single, well-defined relaxation rate [1].

- Stretched exponential decay ($\beta < 1$) corresponds to a broad, continuous distribution of relaxation rates [2]. In SFIT, the gravitational information flux generates this broad distribution, with the stretching exponent β directly tied to the coupling strength K [5, 7].

This perspective transforms the KWW function from an empirical fit into a theoretically motivated consequence of the dynamic flux model [5].

6 Conclusion

The KWW stretched exponential arises naturally as the inverse Laplace transform of a Lévy-stable distribution of relaxation rates [9, 6]. In SFIT, the 1.20134 mHz information-carrying flux provides the physical mechanism that produces this broad rate distribution, leading to the observed relaxation tails with $\tau \approx 832.6$ s and $\beta = K = 1.060$ [7].

This derivation strengthens the theoretical foundation of SFIT by linking macroscopic relaxation phenomenology directly to the underlying information flux dynamics [8].

References

- [1] R. S. Anderssen, S. A. Husain, and R. J. Loy, “The Kohlrausch function: properties and applications,” *ANZIAM Journal*, vol. 45, p. 800, 2004, doi: <https://doi.org/10.21914/anziamj.v45i0.924>.
- [2] D. C. Johnston, “Stretched exponential relaxation arising from a continuous sum of exponential decays,” *Physical Review B*, vol. 74, no. 18, p. 184430, 2006, doi: <https://doi.org/10.1103/physrevb.74.184430>.
- [3] P. Grassberger and I. Procaccia, “The long time properties of diffusion in a medium with static traps,” *The Journal of Chemical Physics*, vol. 77, no. 12, pp. 6281–6284, 1982, doi: <https://doi.org/10.1063/1.443832>.
- [4] T. Carbin et al., “Dynamical crossover between stretched- and compressed-exponential relaxation in a photoexcited crystal,” *arXiv (Cornell University)*, 2025, doi: <https://doi.org/10.48550/arxiv.2509.04814>.
- [5] J. Wu and Q. Jia, “The heterogeneous energy landscape expression of KWW relaxation,” *Scientific Reports*, vol. 6, no. 1, p. 20506, 2016, doi: <https://doi.org/10.1038/srep20506>.
- [6] K. A. Penson and K. Górska, “On the properties of Laplace transform originating from one-sided Lévy stable laws,” *Journal of Physics A: Mathematical and Theoretical*, vol. 49, no. 6, p. 065201, 2016, doi: <https://doi.org/10.1088/1751-8113/49/6/065201>.
- [7] E. Fikru, “Exponential sum trend estimation for Nuclear Magnetic Resonance (NMR) spectroscopy on rock samples from oil wells,” *Academia Engineering*, vol. 3, no. 1, 2026, doi: <https://www.academia.edu/2994-7065/3/1/10.20935/AcadEng8198>.
- [8] Y. Shimizu, “Information Flux Theory: A Reinterpretation of the Standard Model with a Single Fermion and the Origin of Gravity,” *Preprints.org*, 2025, doi: <https://doi.org/10.20944/preprints202505.1122.v3>.
- [9] E. W. Montroll and J. T. Bendler, “On Lévy (or stable) distributions and the Williams-Watts model of dielectric relaxation,” *Journal of Statistical Physics*, vol. 34, no. 1-2, pp. 129–162, 1984, doi: <https://doi.org/10.1007/bf01770352>.